

General Treatment of Orbiting Gyroscope Precession

Ronald J. Adler¹ and Alexander S. Silbergleit¹

Received December 13, 1999

We review the derivation of the metric for a spinning body of any shape and composition using linearized general relativity theory (LGRT), and also obtain the same metric using a transformation argument. The latter derivation makes it clear that the linearized metric contains only the Eddington α and γ parameters, so no new parameter is involved in frame-dragging or Lense–Thirring effects. We then calculate the precession of an orbiting gyroscope in a general weak gravitational field described by a Newtonian potential (the gravitoelectric field) and a vector potential (the gravitomagnetic field). Next we make a multipole analysis of the potentials and the precession equations, giving all of these in terms of the spherical harmonics moments of the density distribution. The analysis is not limited to an axially symmetric source, although the Earth, which is the main application, is very nearly axisymmetric. Finally, we analyze the precession in regard to the Gravity Probe B (GP-B) experiment, and find that the effect of the Earth's quadrupole moment (J_2) on the geodetic precession is large enough to be measured by GP-B (a previously known result), but the effect on the Lense–Thirring precession is somewhat beyond the expected GP-B accuracy.

1. INTRODUCTION

The Gravity Probe B satellite is scheduled to fly in the year 2000 (Keiser, 1998). It contains a set of gyroscopes intended to test the predictions of general relativity (GR) that a gyroscope in a low circular polar orbit with altitude 650 km will precess about 6.6 arcsec/year in the orbital plane (geodetic precession) and about 42 milliarcsec/year perpendicular to the orbital plane [LT precession, see Misner *et al.* (1973); Ohanian and Ruffini (1994), secs. 4.7 and 7.8; Will (1993), sec. 9.1]. In this paper we review the theoretical derivation of these effects and in particular consider the contributions of the Earth's quadrupole and higher multipole fields.

¹Gravity Probe B, W. W. Hansen Experimental Physics Laboratory, Stanford University, Stanford, California 94305-4085; e-mail: adler@relgyro.stanford.edu, gleit@relgyro.stanford.edu

We first review the derivation of the metric for a rotating body using the standard LGRT approach (Lense and Thirring, 1918; Misner *et al.*, 1973; Ohanian and Ruffini, 1994, secs. 4.7 and 7.8). The metric is characterized by a Newtonian scalar potential (the gravitoelectric field) and a vector potential (the gravitomagnetic field) (Ohanian and Ruffini, 1994, sec. 3.5). We then obtain the same result with a simple transformation argument which clarifies the physical meaning of the metric (Will, 1993, sec. 4.3). Specifically, it makes clear that if the metric of a point mass contains fundamental parameters such as the Eddington parameters α and γ , then to lowest order the metric of a rotating body contains no *new* fundamental parameters (Nordtvedt, 1998). Thus there is no new Lense–Thirring or frame-dragging parameter to be measured by GP-B or any other experiment.

We then derive the precession equations for a gyroscope in a general weak field system, that is, for any scalar and vector potential fields (Misner *et al.*, Wheeler, 1973). The calculation is valid to first order in the fields and velocities of the source body and the satellite. The gravitational field of the Earth is described by the scalar and vector potentials, which depend on the shape of the body and the mass and velocity distribution inside it. We treat both of these fields by a multipole expansion and express the precessions in series of spherical harmonics. We do not limit ourselves to the axially symmetric case, which has been thoroughly studied by Teyssandier (1977, 1978). In particular, we show that for a solid body rotation up to the order $l \leq 2$ both precessions depend only on the tensor of inertia of the Earth.

The major nonspherical contributions to the GP-B precessions are from the Earth's quadrupole moment. The contribution to the geodetic precession which has a magnitude of about 1 part in 10^3 is detectable by GP-B and quite important for the determination of the parameter γ , which is to be measured to about 1 part in 10^5 , the most accurate measurement envisioned (Ohanian and Ruffini, 1994, sec. 3.5 and in particular Table 14.2). This contribution has been calculated independently by Wilkins (1968) and Barker and O'Connell (1970), and then in the most elegant and general way by Breakwell (1988). The contribution to the Lense–Thirring precession is beyond the expected accuracy of GP-B and close enough to the result by Teyssandier (1977) obtained from a different Earth model.

We also estimate the influence of Moon and Sun to find that only the geodetic contribution of the Sun must be included in the GP-B data reduction, as anticipated.

Since many parts of this paper refer to previously known results, a brief summary of new and novel features is in order. The derivation of the Lense–Thirring metric by Lorentz transformation and superposition is the simplest and most physical analysis of which we are aware; the physical conclusion is that no independent new Eddington (or PPN) parameter is

associated with gravitomagnetic effects, which has involved some dispute (Ciufolini and Wheeler, 1995). The basic equations for the precession are derived in complete generality in the context of the linearized theory, and the “small” terms are considered in detail. The multipole expansions are for general (quasistatic) scalar and vector fields and not limited to axisymmetric fields; this allows for future analysis of nonaxisymmetric perturbations for GP-B and astrophysical systems. For the case where only the multipole moments for $e \leq 2$ are important, particularly simple precession results are given in terms of the inertia tensor; this allows extension of the classic Lense–Thirring formula to a spherical body with any density distribution, or to a symmetric top rotating about its symmetry axis. Using two rather general models of the Earth, we illustrate how the vector potential field may be constructed from the known scalar potential using our general relations. Finally, the oblateness correction to the Lense–Thirring precession confirms the result of Teysandier that it is small enough that it does not need to be included in the GP-B analysis.

2. THE METRIC IN LINEARIZED GENERAL RELATIVITY THEORY

We first briefly review this standard derivation of linearized general relativity theory. The metric of a rotating body such as the Earth is obtained by introducing a small perturbation of the Lorentz metric $\eta_{\mu\nu}$, that is,

$$g_{\mu\nu} = \eta_{\mu\nu} + h_{\mu\nu} \quad (1)$$

The perturbation $h_{\mu\nu}$ is assumed to be expressed in isotropic space coordinates so that $h_{11} = h_{22} = h_{33} = h_s$. Similarly, we suppose the matter producing the metric field is described by the energy-momentum tensor of slow-moving and low-density matter with negligible pressure,

$$T^{\mu\nu} = \rho u^\mu u^\nu \quad (2)$$

where u^μ is the 4-velocity and ρ is the matter density. The field equations are

$$G_{\mu\nu} = 8\pi G T_{\mu\nu} \quad (3)$$

where $G_{\mu\nu}$ is the Einstein tensor.

The calculation of $G_{\mu\nu}$ and $T_{\mu\nu}$ to the lowest order in the perturbation is straightforward and results in the following equations:

$$D^2 \left(h_{\mu\nu} - \frac{1}{2} h \eta_{\mu\nu} \right) = 16\pi G T_{\mu\nu}, \quad h \equiv h^\sigma_\sigma, \quad D^2 \equiv \frac{\partial^2}{\partial t^2} - \nabla^2 \quad (4)$$

$$h^\mu{}_\nu = 0 \quad (5)$$

Here indices are raised and lowered with the Lorentz metric, which is consistent to lowest order, and the vertical bar denotes differentiation. The last expression imposes the so-called Lorentz condition, which can always be achieved by a coordinate transformation (to lowest order) and involves no loss of generality; it is also the analog of a gauge choice in electromagnetism. Because we have used isotropic coordinates, equation (4) leads immediately to $h_{00} = h_s$, and the standard classical correspondence gives $h_{00} = h_s = 2\Phi$. We thereby obtain the wave equations for the scalar and vector potentials:

$$D^2\Phi = -4\pi G\rho \quad (6)$$

$$D^2\vec{h} = 16\pi G\rho\vec{v}, \quad \vec{h} = \{h_{01}, h_{02}, h_{03}\} \quad (7)$$

where \vec{v} is the velocity of the source. These equations may be solved in the standard way by means of a retarded Green's function. In the case of a time-independent system, or a system which changes so slowly that retardation effects may be ignored, the solution is

$$\Phi(\vec{r}) = -G \int \frac{\rho(\vec{r}') d^3r'}{|\vec{r} - \vec{r}'|} \quad (8)$$

$$\vec{h}(\vec{r}) = 4G \int \frac{\rho(\vec{r}')\vec{v}(\vec{r}') d^3r'}{|\vec{r} - \vec{r}'|} \quad (9)$$

Note that these expressions are analogs of the equations of electrostatics and magnetostatics, which is why one may speak about gravitoelectric and gravitomagnetic effects in the linearized theory.

In summary, we may write the (Lense-Thirring) line element as

$$ds^2 = (1 + 2\Phi) dt^2 - (1 - 2\Phi) d\vec{r}^2 + 2\vec{h} \cdot d\vec{r} dt \quad (10)$$

This is valid to first order in field intensity and source velocity, and will serve as a basis for calculating the gyroscope precession. Figure 1 shows the general aspects of the scalar and vector fields of a spinning body; the vector field generally points in the same direction as the velocity of the surface of the body.

3. THE METRIC WITH PARAMETERS: DERIVATION BY FRAME TRANSFORMATION

It is, in fact, possible to obtain the above result from a different and physically interesting perspective, and moreover introduce parameters convenient for discussing experimental measurements. Following Eddington, consider the metric of a massive point with a geometric mass m at a large distance r , so that $m/r \ll 1$. We expand the Schwarzschild solution in isotropic coordinates for this situation as




Fig. 1. (a) The scalar field contours and the flow lines of the vector field. (b) The gradient of the scalar field and the curl of the vector field.

$$\begin{aligned}
 ds^2 &= \frac{(1 - m/2r)^2}{(1 + m/2r)^2} dt^2 - \left(1 - \frac{m}{2r}\right)^4 d\vec{r}^2 \\
 &= \left(1 - \frac{2m}{r} + \frac{2m^2}{r^2} + \dots\right) dt^2 - \left(1 + \frac{2m}{r} + \frac{3m^2}{r^2} + \dots\right) d\vec{r}^2
 \end{aligned}$$

Eddington suggested that this be written in terms of parameters as

$$ds^2 = \left(1 - \alpha \frac{2m}{r} + \beta \frac{2m^2}{r^2} + \dots\right) dt^2 - \left(1 + \gamma \frac{2m}{r} + \dots\right) d\vec{r}^2 \quad (11)$$

where α , β , and γ are equal to 1 for general relativity. The power series (11) is clearly a rather general form for the metric far from a spherical body.

Since the parameter m which appears in the metric is a constant of integration representing the mass of the central body (specifically $m = GM/c^2$), we may absorb the parameter α into it, which is equivalent to taking $\alpha \equiv 1$. This is consistent as long as no independent nongravitational determination of the mass of the body is considered. We will nevertheless retain α in our calculations as a book-keeping device.

Indeed, the parameters in (11) may all be viewed as a tool for tracking which terms in the metric contribute to a gravitational phenomenon such as the perihelion shift of Mercury or the deflection of starlight by the Sun. Alternatively they may be viewed as numbers which may *not* be equal to 1 if a metric theory other than general relativity is actually valid. In either case they provide a convenient way to express the results of experimental tests of gravity as giving value to the parameters. This parametrized approach has

been extended to include many other parameters and has been highly developed under the name "parametrized post-Newtonian theory," or PPN (Will, 1993). In this paper we take the viewpoint that general relativity is to be tested and emphasize that we are *not* using the more general PPN approach. Solar system measurements give $\beta - 1 = (0.2 \pm 1.0) \times 10^{-3}$ and $\gamma - 1 = (-1.2 \pm 1.6) \times 10^{-3}$, which are of course entirely consistent with general relativity.

We consider now only phenomena in which the quadratic term, $\sim m^2/r^2$, in g_{00} is unimportant, that is, in which we may ignore β and assume that the underlying gravitational theory is linear. Then for a stationary mass,

$$ds^2 = \left(1 - \alpha \frac{2m}{r}\right) dt^2 - \left(1 + \gamma \frac{2m}{r}\right) d\vec{r}^2$$

(stationary point mass) (12)

Since this is nearly the Lorentz metric, we may generalize it to a moving mass point by simply transforming to a moving system using a transformation that is Lorentzian to the first order in velocity:

$$t_r = t - vx, \quad x_r = x - vt$$

Here the subscript r labels the system in which the mass is at rest, and which moves at velocity v in the x direction in the laboratory system. This transformation gives the metric for the moving mass point as

$$ds^2 = \left(1 - \alpha \frac{2m}{r}\right) dt^2 - \left(1 + \gamma \frac{2m}{r}\right) d\vec{r}^2 + (\alpha + \gamma) \frac{4m}{r} v dx dt$$

(point mass moving in x direction)

This obviously generalizes for motion in any direction to

$$ds^2 = \left(1 - \alpha \frac{2m}{r}\right) dt^2 - \left(1 + \gamma \frac{2m}{r}\right) d\vec{r}^2 + (\alpha + \gamma) \frac{4m}{r} (\vec{v} \cdot d\vec{r}) dt$$

(moving point mass) (13)

As we assume that our theory is linear to this order (like general relativity), we can superpose the fields of a distribution of such point masses and replace m/r by $\Phi(\vec{r})$ from (8) and $4m\vec{v}/r$ by $\vec{h}(\vec{r})$ from (9), resulting in

$$\Phi(\vec{r}) = -G \int \frac{\rho(\vec{r}') d^3\vec{r}'}{|\vec{r} - \vec{r}'|}$$

$$\vec{h}(\vec{r}) = 4G \int \frac{\rho(\vec{r}') \vec{v}(\vec{r}') d^3\vec{r}'}{|\vec{r} - \vec{r}'|}$$

$$ds^2 = (1 + 2\alpha\Phi) dt^2 - (1 - 2\gamma\Phi) d\vec{r}^2 + (\alpha + \gamma)(\vec{h} \cdot d\vec{r}) dt \quad (14)$$

This agrees with the general relativity result (10) when $\alpha = \gamma = 1$, but now contains appropriate combinations of Eddington parameters.

We emphasize that this line element has been obtained for any slowly moving mass distribution from the parametrized metric for a stationary mass point by transformation and superposition, and thus no new parameter appears in the expression. Therefore a measurement of a phenomenon which depends on the cross term in (14) provides a value for $\alpha + \gamma$ and *not* for some new parameter associated with gravitomagnetism (Cuifollini and Wheeler, 1995; Nordtvedt, 1968, 1988a, b, 1998; Adler and Silbergleit, 1998; Adler, 1999).

Note also that the result is rather strong since it depends on the observationally verified Schwarzschild metric (12), the well-tested Lorenz transformation, and the principle of superposition, valid for any linear theory.

4. GENERAL PRECESSION RELATIONS

An orbiting gyroscope has its spin axis parallel-displaced in accord with the metric (14). We calculate this motion with minimal assumptions about the potentials, which could be the potentials of a nearly spherical and rather uniform rigid body such as the Earth or a potato-shaped body such as in Fig. 1, with some interior mass distribution. We will work always to the first order in the potentials and in the velocities \vec{v} and \vec{V} of the source and orbiting gyroscope. The parallel displacement equation for the gyro spin S^μ is (Misner *et al.*, 1973; Ohanian and Ruffini, 1994, secs. 4.7 and 7.8; Will, 1993, sec. 9.1)

$$\frac{dS^\mu}{ds} + \left\{ \begin{matrix} \mu \\ \nu\sigma \end{matrix} \right\} S^\nu \frac{dx^\sigma}{ds} = 0 \quad (15)$$

We suppose that the gyro spin 4-vector is perpendicular to the velocity 4-vector, which is equivalent to assuming that the gyro spin has no zero component in its rest frame. From this assumption the zero component in another frame is easily obtained, and to first order in the satellite velocity it is given by $S^0 = \vec{S} \cdot \vec{V}$. We calculate the Euler–Lagrange equations in the standard way, and then put them into canonical form to give the Christoffel symbols. To lowest order in the potentials and velocity the Christoffel symbols are

$$\begin{aligned} \left\{ \begin{matrix} 0 \\ 0l \end{matrix} \right\} &= \alpha\Phi_{|l}, & \left\{ \begin{matrix} i \\ 00 \end{matrix} \right\} &= \alpha\Phi_{|i}, & \left\{ \begin{matrix} i \\ ii \end{matrix} \right\} &= -\gamma\Phi_{|i} \\ \left\{ \begin{matrix} i \\ il \end{matrix} \right\} &= -\gamma\Phi_{|l}, & \left\{ \begin{matrix} i \\ ll \end{matrix} \right\} &= \gamma\Phi_{|i}, & \left\{ \begin{matrix} i \\ 0l \end{matrix} \right\} &= \frac{\alpha + \gamma}{4} (h_{|li} - h_{|il}) \end{aligned}$$

The Roman indices in these expressions are space indices and run from 1 to 3, a vertical bar denotes an ordinary derivative, and $l \neq i$. Note that the gravitational vector potential \vec{h} occurs in a gauge invariant way, that is only its *curl* appears.

Substitution of the Christoffel symbols into the spin equation of motion gives

$$\begin{aligned} \dot{S}^i + (\alpha + \gamma)\Phi_{|i}(S^l V^l) - \gamma S^i(\Phi_{|l} V^l) - \gamma V^i(\Phi_{|l} S^l) \\ + \frac{\alpha + \gamma}{4}(h_{|i} - h_{|l})S^l = 0 \end{aligned} \quad (16)$$

We break the drift rate into two parts, the geodetic drift rate due to the scalar potential Φ , and the Lense–Thirring precession rate due to the vector potential \vec{h} . Separating also symmetric and antisymmetric parts of the geodetic effect, we arrive at, in a 3-dimensional vector notation,

$$\dot{\vec{S}}_{LT} = \vec{\Omega}_{LT} \times \vec{S}, \quad \vec{\Omega}_{LT} \equiv \left(\frac{\alpha + \gamma}{4}\right) \nabla \times \vec{h} \quad (17)$$

and

$$\begin{aligned} \dot{\vec{S}}_G = \vec{\Omega}_G \times \vec{S} + \left\{ -\frac{\alpha}{2} [(\vec{S} \cdot \vec{V})\nabla\Phi + (\vec{S} \cdot \nabla\Phi)\vec{V}] + \gamma(\vec{V} \cdot \nabla\Phi)\vec{S} \right\} \\ \vec{\Omega}_G \equiv \left(\frac{\alpha + 2\gamma}{2}\right) \nabla\Phi \times \vec{V} \end{aligned} \quad (18)$$

$$\dot{\vec{S}} = \dot{\vec{S}}_G + \dot{\vec{S}}_{LT} \quad (19)$$

where $\vec{\Omega}_{LT}$ and $\vec{\Omega}_G$ are the instantaneous values of the Lense–Thirring and geodetic precessions, respectively. Since $\vec{\Omega}_{LT}$ is the *curl* of the gravitational vector potential the Lense–Thirring precession rate is the analog of the magnetic field in magnetostatics theory. The geodetic effect is of the order of the scalar potential times the orbital velocity of the satellite, while the Lense–Thirring terms are of the order the scalar potential times the velocity of the central body (the Earth), which is typically much smaller than the orbital velocity. Symmetric geodetic terms are responsible for stretching the spin vector \vec{S} , and the reason for separating them is that their effect almost vanishes when averaged over any reasonable satellite orbit. To see this, consider the last term in the symmetric part of the geodetic drift rate. Using Newton's law in the form $\nabla\Phi = -\vec{V}$, we may write

$$\langle (\nabla\Phi \cdot \vec{V}) \rangle \vec{S} = - \left\langle \left(\frac{d\vec{V}}{dt} \cdot \vec{V} \right) \right\rangle \vec{S} = - \left\langle \frac{1}{2} \frac{d\vec{V}^2}{dt} \right\rangle \vec{S} = - \frac{\Delta\vec{V}^2}{2T} \vec{S}$$

where $\Delta\vec{V}^2$ is the change in the velocity squared in total time T , and it is assumed that the drift rate is small and \vec{S} does not change significantly during this period of time. If the orbit is periodic, this quantity will be zero, and for a nearly periodic orbit it will be very small. Similarly for the remaining symmetric terms in the geodetic drift we may write

$$\begin{aligned} \langle (\vec{S} \cdot \vec{V}) \nabla\Phi + (\vec{S} \cdot \nabla\Phi)\vec{V} \rangle &= - \left\langle (\vec{S} \cdot \vec{V}) \frac{d\vec{V}}{dt} + \left(\vec{S} \cdot \frac{d\vec{V}}{dt} \right) \vec{V} \right\rangle \\ &= - \left\langle V^i \frac{dV^i}{dt} + V^i \frac{dV^i}{dt} \right\rangle S^i \\ &= - \left\langle \frac{dV^i V^i}{dt} \right\rangle S^i = - \frac{\Delta V^i V^i}{T} S^i \end{aligned}$$

Again, this is zero for a periodic orbit and very small for a nearly periodic orbit.

In summary, the average precession rate of the gyro spin is

$$\langle \vec{S} \rangle = \langle \vec{S}_G \rangle + \langle \vec{S}_{LT} \rangle, \quad \langle \vec{S}_G \rangle = \langle \vec{\Omega}_G \rangle \times \vec{S}, \quad \langle \vec{S}_{LT} \rangle = \langle \vec{\Omega}_{LT} \rangle \times \vec{S} \quad (20)$$

with the values of the geodetic and precessions given in (17) and (18).

Note that the geodetic and Lense–Thirring effects are approximately perpendicular to each other for an approximately polar orbit around a central body of reasonable shape; for a circular polar orbit about a spherical body they are perpendicular. Figure 2 shows the orientation of the various vectors for a fairly general situation, a satellite in a roughly planar polar orbit about an oddly shaped body. The geodetic precession vector $\vec{\Omega}_G$ is roughly perpendicular to the orbit plane and the Lense–Thirring precession vector $\vec{\Omega}_{LT}$ has the general appearance of a dipole magnetic field.

5. EFFECT OF DISTANT MASSES

If there are relatively distant masses, such as the Moon, in the neighborhood of the central body, their density distribution $\rho_D(\vec{r})$ may be expressed as

$$\rho_D(\vec{r}) = \sum_n M_n \delta(\vec{r} - \vec{r}_n)$$

The effect on the scalar and vector potentials is

$$\Phi_D(\vec{r}) = - \sum_n \frac{GM_n}{|\vec{r} - \vec{r}_n|}, \quad \vec{h}_D(\vec{r}) = 4 \sum_n \frac{GM_n \vec{v}_n}{|\vec{r} - \vec{r}_n|}$$

It then follows that the precessions due to these distant masses are




Fig. 2. (a) The satellite is in a roughly planar polar orbit, and the geodetic precession vector is perpendicular to the orbit. (b) The orientation of the vectors relevant to the Lense-Thirring precession: the precession vector field has the appearance of a dipole magnetic field.

$$\vec{\Omega}_G^D = -\frac{\alpha + 2\gamma}{2} \sum_n \frac{GM_n(\vec{r} - \vec{r}_n) \times \vec{V}}{|\vec{r} - \vec{r}_n|^3}, \quad \vec{\Omega}_{LT}^D = -(\alpha + \gamma) \sum_n \frac{G\vec{L}_n}{|\vec{r} - \vec{r}_n|^3} \tag{21}$$

where $\vec{L}_n = M_n(\vec{r} - \vec{r}_n) \times \vec{v}_n$ is the angular momentum of the distant mass.

For the specific case of the Moon and the GP-B experiment the numerical values will be discussed in Section 9.

6. MULTIPOLE EXPANSIONS

We will henceforth study the case of a body such as the Earth which is rigidly rotating, that is, $\vec{v} = \vec{\omega} \times \vec{r}$. Motivated by expression (9), we introduce a new vector potential quantity $\vec{\Pi}(\vec{r})$:

$$\vec{h}(\vec{r}) = 4\vec{\omega} \times \vec{\Pi}(\vec{r}), \quad \vec{\Pi}(\vec{r}) = G \int \frac{\rho(\vec{r}')\vec{r}' d^3r'}{|\vec{r} - \vec{r}'|} \tag{22}$$

The vector $\vec{\Pi}(\vec{r})$ is a harmonic function outside the body which we will expand in spherical harmonics; for a special spherically symmetric case $\vec{\Pi}(\vec{r})$ is collinear with \vec{r} . In terms of $\vec{\Pi}(\vec{r})$, the metric (14) may be written as

$$ds^2 = (1 + 2\alpha\Phi) dt^2 - (1 - 2\gamma\Phi) d\vec{r}^2 + 4(\alpha + \gamma) (\vec{\omega} \times \vec{\Pi}) \cdot d\vec{r} dt \tag{23}$$

Using (8) and (9), it is possible to express the divergence of $\vec{\Pi}(\vec{r})$ via the scalar potential, namely,

$$\nabla \cdot \vec{\Pi}' = -(\Phi + \vec{r} \cdot \nabla\Phi)$$

which allows us to rewrite formula (17) for the Lense–Thirring precession as

$$\begin{aligned} \vec{\Omega}_{LT} &= -(\alpha + \gamma) [(\vec{\omega} \cdot \nabla)\vec{\Pi} + \vec{\omega}(\Phi + \vec{r} \cdot \nabla\Phi)] \vec{\Omega}_{LT} \\ &= -(\alpha + \gamma)\omega [\vec{\Pi}'_{|z} + (\Phi + r\Phi_{|r})\hat{z}] \end{aligned} \tag{24}$$

Here we have chosen the z axis to be along the spin, $\vec{\omega} = \omega\hat{z}$.

We now put the origin of spherical coordinates $\{r, \theta, \varphi\}$ at the center of mass of the body, introduce spherical harmonics with the notation

$$Y_{lm}^v(\theta, \varphi) = P_l^m(\cos \theta)f^v(m\varphi), \quad f^v(m\varphi) = \begin{cases} \cos m\varphi, & v = c \\ \sin m\varphi, & v = s \end{cases} \tag{25}$$

and expand the potentials Φ and $\vec{\Pi}$ in corresponding series:

$$\Phi(\vec{r}, t) = -\frac{GM}{r} \left[1 + \sum_{l \geq 2, m, v} a_{lm}^v(t) \left(\frac{R}{r}\right)^l Y_{lm}^v(\theta, \varphi) \right] \tag{26}$$

$$\Pi_i(\vec{r}, t) = \frac{GMR}{r} \left[\sum_{l \geq 1, m, v} p_{lm}^{iv}(t) \left(\frac{R}{r}\right)^l Y_{lm}^v(\theta, \varphi) \right] \tag{27}$$

[$l = 1$ terms in (26) and $l = 0$ terms in (27) are missing because the origin is at the center of mass of the body; see expressions (28) and (29) below]. These expansions are in the inertial frame in which the body rotates; the direction of the coordinate polar axis z is so far arbitrary; R is the characteristic size of the body, which we take to be the equatorial radius for the Earth. The potentials are slowly varying functions of time due to the Earth rotation, so we write the coefficients as explicit functions of time. These coefficients are related to the mass distribution in the standard way by

$$a_{lm}^v(t) = \frac{(2 - \delta_{m0})(l - m)!}{M(l + m)!} \int \rho(\vec{r}, t) \left(\frac{r}{R}\right)^l Y_{lm}^v(\theta, \varphi) d^3\vec{r} \tag{28}$$

$$p_{lm}^{iv}(t) = \frac{(2 - \delta_{m0})(l - m)!}{M(l + m)!} \int \rho(\vec{r}, t) \left(\frac{r}{R}\right)^l \frac{x_i}{R} Y_{lm}^v(\theta, \varphi) d^3\vec{r} \tag{29}$$

Note that generally the domain of integration here is also time dependent. In particular, we write for convenience $a_{i0}^c(t) \equiv a_{i0}(t)$, $a_{i0}^s(t) \equiv 0$, $p_{i0}^c(t) \equiv p_{i0}^c(t)$, $p_{i0}^s(t) \equiv 0$.

The time dependence in (28) and (29) may be easily analyzed for a rigidly rotating body provided that the rotation axis is fixed both in the inertial space and in the body. In that case in the inertial frame with the z axis along the spin, $\vec{\omega} = \omega\hat{z}$, the Earth density is

$$\rho(\vec{r}, t) = \rho_e(r, \theta, \varphi - \omega t) \equiv \rho_e(r, \theta, \bar{\varphi})$$

where $\rho_e(r, \theta, \bar{\varphi}) \equiv \rho_e(\vec{r})$ is the time-independent density measured in the frame rotating with the Earth. Substituting this for the density in (28) and (29) and transforming to the Earth-fixed frame, we find that the time-dependent coefficients may be written in terms of constant moments of density according to

$$\begin{bmatrix} a_{lm}^c(t) \\ a_{lm}^s(t) \end{bmatrix} = \begin{bmatrix} \cos \omega t & -\sin \omega t \\ \sin \omega t & \cos \omega t \end{bmatrix} \begin{bmatrix} a_{lm}^c \\ a_{lm}^s \end{bmatrix} \tag{30}$$

$$\begin{bmatrix} p_{lm}^{ic}(t) \\ p_{lm}^{is}(t) \end{bmatrix} = \begin{bmatrix} \cos \omega t & -\sin \omega t \\ \sin \omega t & \cos \omega t \end{bmatrix} \begin{bmatrix} p_{lm}^{ic} \\ p_{lm}^{is} \end{bmatrix} \tag{31}$$

and the time-independent coefficients are explicitly given by

$$a_{lm}^v = a_{lm}^v(0) = \frac{(2 - \delta_{m0})(l - m)!}{M(l + m)!} \int \rho_e(\vec{r}) \left(\frac{r}{R}\right)^l Y_{lm}^v(\theta, \bar{\varphi}) d^3\vec{r} \tag{32}$$

$$p_{lm}^{iv} = p_{lm}^{iv}(0) = \frac{(2 - \delta_{m0})(l - m)!}{M(l + m)!} \int \rho_e(\vec{r}) \left(\frac{r}{R}\right)^l \frac{x_i}{R} Y_{lm}^v(\theta, \bar{\varphi}) d^3\vec{r} \tag{33}$$

If the rotation axis wanders in the body and/or in the inertial space, the relation between the time-dependent and time-independent coefficients is given by a combination of appropriate rotation matrices which is more complicated than the one matrix in (30) and (31) (Rose, 1957). Generally, a time-dependent coefficient with the indices l and m is a linear combination of the appropriate time-independent coefficients with the indices l and $n = 0, 1, \dots, l$.

Introducing now the time-dependent coefficients from (32) and (33) back into (26) and (27), we obtain the time-dependent potentials in a convenient form using time-independent coefficients only:

$$\Phi(\vec{r}, t) = -\frac{GM}{r} \left[1 + \sum_{l \geq 2, m, v} a_{lm}^v \left(\frac{R}{r}\right)^l Y_{lm}^v(\theta, \varphi - \omega t) \right] \tag{34}$$

$$\Pi_i(\vec{r}, t) = \frac{GMR}{r} \left[\sum_{l \geq 1, m, v} p_{lm}^{iv} \left(\frac{R}{r}\right)^l Y_{lm}^v(\theta, \varphi - \omega t) \right] \tag{35}$$

This is of course what one should expect intuitively; in general this form will be most useful for our purposes.

The constant coefficients a_{lm}^v in (34) are those that are measured very accurately for the Earth [up to $l = 18$, their values are found in *World Geodetic System 1984* (1987)]. However, for the Earth of arbitrary shape and

composition it is impossible to express p_{lm}^{iv} through a_{lm}^v ; in other words, their values, and hence the scalar and vector potentials, are independent. Nevertheless, a useful relationship between the two sets of coefficients exists; to describe it, we need a notation for a general moment of the density,

$$M_{klm}^v \equiv \int \rho(\vec{r}) \left(\frac{r}{R}\right)^k Y_{lm}^v(\theta, \varphi) d^3r \tag{36}$$

In particular,

$$a_{lm}^v = \frac{(2 - \delta_{m0})(l - m)!}{M(l + m)!} M_{ilm}^v \tag{37}$$

Using the definitions (32) and (33) and the recurrence relations for Legendre functions (Bateman and Erdélyi, 1953), we derive the following equalities relating p_{lm}^{iv} to a_{lm}^v :

$$\begin{aligned} p_{lm}^{1v} &= (2l + 1)^{-1} \{ -2^{-1}(l + m + 1)(l + m + 2)a_{l+1m+1}^v \\ &\quad + (2 - \delta_{m1})^{-1}a_{l+1m-1}^v + [(l - m)!/M(l + m)!] \\ &\quad [M_{l+1l-1m+1}^v - (l + m - 1)(l + m)M_{l+1l-1m-1}^v] \} \\ p_{lm}^{2v} &= (\mp)(2l + 1)^{-1} \{ 2^{-1}(l + m + 1)(l + m + 2)a_{l+1m+1}^\mu \\ &\quad + (2 - \delta_{m1})^{-1}a_{l+1m-1}^\mu - [(l - m)!/M(l + m)!] \\ &\quad [M_{l+1l-1m+1}^\mu - (l + m - 1)(l + m)M_{l+1l-1m-1}^\mu] \} \\ p_{lm}^{3v} &= (2l + 1)^{-1} \{ (l + m + 1)a_{l+1m+1}^v \\ &\quad + (2 - \delta_{m0})(l - m)!M_{l+1l-1m}^v/M(l + m - 1)! \} \end{aligned} \tag{38}$$

In the second line of (38), the minus sign is taken and $\mu = s$ when $\nu = c$, the plus sign and $\mu = c$ when $\nu = s$.

From (34) and (35), using the definition (17) of $\vec{\Omega}_{LT}$, we can compute a rather compact multipole expansion for the Lense–Thirring precession,

$$\begin{aligned} \Omega_{LT}^i(\vec{r}, t) &= \frac{\alpha + \gamma}{2} \frac{GM\omega}{r} \sum_{l \geq 2, m, \nu} [(l - m)p_{l-\delta_{l3m}}^{iv} \\ &\quad - la_{lm}^v \delta_{i3}] \left(\frac{R}{r}\right)^l Y_{lm}^v(\theta, \varphi - \omega t) \end{aligned} \tag{39}$$

The corresponding expansion for the geodetic precession is too cumbersome to be useful.

7. THE FAR-FIELD/HIGH-SYMMETRY APPROXIMATION: MULTIPOLES WITH $l \leq 2$ AND THE TENSOR OF INERTIA

If the shape of the central body and the mass distribution inside it are known, then all the pertinent quantities, including multipole expansion coefficients a_{lm}^v and p_{lm}^{iv} , may be found by integration, but this is rarely the case. Even when a_{lm}^v are measured, as for the Earth, all the p_{lm}^{iv} , and the Lense-Thirring effect with them, remain entirely undetermined. However, for a body of any shape and composition, the coefficients a_{2m}^v ($l = 2$) and p_{1m}^{iv} ($l = 1$) can be expressed in terms of the elements I_{ij} of the tensor of inertia \mathbf{I} determined in a standard way,

$$I_{ij} = \int \rho(\vec{r})(r^2 \delta_{ij} - x_i x_j) d^3\vec{r}$$

Writing $I_{ii} \equiv I_i$, we find

$$\begin{aligned} a_{20} &= -\frac{2I_3 - I_2 - I_1}{2MR^2}, & a_{22}^c &= \frac{I_2 - I_1}{4MR^2}, & p_{10}^3 &= -\frac{I_3 - I_2 - I_1}{2MR^2} \\ p_{11}^{1c} &= -\frac{I_3 + I_2 - I_1}{2MR^2}, & p_{11}^{2s} &= -\frac{I_3 - I_2 + I_1}{2MR^2} \\ a_{21}^c &= -p_{10}^1 = p_{11}^{3c} = \frac{I_{13}}{MR^2}, & a_{21}^s &= -p_{10}^2 = p_{11}^{3s} = \frac{I_{23}}{MR^2} \\ -a_{22}^s &= 0.5 p_{11}^{1s} = 0.5 p_{11}^{2c} = \frac{I_{12}}{MR^2} \end{aligned} \quad (40)$$

This is done by comparing the integrals (32) (with $l = 2$) and (33) (with $l = 1$) to I_{ij} using explicit expressions of Legendre functions with $l = 1, 2$; formulas for a_{20} and a_{22}^c are known and used in geodesy for the determination of the Earth's moments of inertia (Bursa, 1992).

Introducing (40) into (34), (35) and dropping the terms with $l > 2$ for Φ and $l > 1$ for $\bar{\Pi}$, we first obtain the $l \leq 2$ formulas for the potentials:

$$\begin{aligned} \Phi(\vec{r}) &= -\frac{G}{r} \left[M + \frac{1}{2r^2} \left(\text{tr } \mathbf{I} - \frac{3}{r^2} (\mathbf{I}\vec{r} \cdot \vec{r}) \right) \right], \\ \bar{\Pi}(\vec{r}) &= -\frac{G}{r^3} \left[\mathbf{I}\vec{r} - \frac{1}{2} (\text{tr } \mathbf{I})\vec{r} \right] \end{aligned} \quad (41)$$

From these we then obtain the approximation for the precessions by differentiation of the expressions (18) and (17) [see also (22)]:

$$\begin{aligned} \vec{\Omega}_G = \frac{(\alpha + 2\gamma)G}{2r^3} \left\{ \left[M + \frac{3}{2r^2} \left(\text{tr } \mathbf{I} - \frac{5}{r^2} (\mathbf{I}\vec{r} \cdot \vec{r}) \right) \right] (\vec{r} \times \vec{V}) \right. \\ \left. + \frac{3}{r^2} (\mathbf{I}\vec{r} \times \vec{V}) \right\} \end{aligned} \tag{42}$$

$$\begin{aligned} \vec{\Omega}_{LT} = \frac{(\alpha + \gamma)G}{r^3} \left\{ \mathbf{I}\vec{\omega} - 3 \left[\frac{1}{2} (\text{tr } \mathbf{I}) - \frac{1}{r^2} (\mathbf{I}\vec{r} \cdot \vec{r}) \right] \vec{\omega} \right. \\ \left. + 3 \frac{\vec{\omega} \cdot \vec{r}}{r^2} \left[\frac{1}{2} (\text{tr } \mathbf{I})\vec{r} - \mathbf{I}\vec{r} \right] \right\} \end{aligned} \tag{43}$$

[Of course, the same expression for $\vec{\Omega}_{LT}$ is also obtained from (40) and (39) with $l = 2$.]

Note that formulas (40) and (41) hold in both the body-fixed and inertial frames, and the coefficients and moments of inertia in the latter depend on the time if the rotation is not axisymmetric, $a_{lm}^v = a_{lm}^v(t)$, $p_{lm}^v = p_{lm}^v(t)$, $I_{ij} = I_{ij}(t)$. The expressions (42) and (43) for the precessions are meaningful in the inertial frame, where generally $\mathbf{I} = \mathbf{I}(t) = \mathbf{R}(t)\mathbf{I}(0)\mathbf{R}^T(t)$, $\mathbf{R}(t)$ being a rotation matrix which converts the body-fixed radius vector into the inertial one. In the above case of a simple rotation about an inertially fixed axis, $\vec{\omega} = \omega\hat{z}$,

$$\mathbf{R}(t) = \begin{vmatrix} \cos \omega t & -\sin \omega t & 0 \\ \sin \omega t & \cos \omega t & 0 \\ 0 & 0 & 1 \end{vmatrix}$$

The results (41)–(43) are valid under either of the two conditions (1) *far field*, $R/r \ll 1$; (2) *high symmetry*, i.e., all higher order moments are small. These results alter somewhat our usual notion of the geodetic and Lense–Thirring effects: the first one is generally proportional not only to the orbital momentum, but to $\mathbf{I}\vec{r} \times \vec{V}$ as well, and the Lense–Thirring precession generally points not only in the direction of the angular momentum $\vec{L} = \mathbf{I}\vec{\omega}$, but has also components parallel to $\vec{\omega}$, \vec{r} , and $\mathbf{I}\vec{r}$. Two particular cases of the inertia tensor are of special interest.

(a) *Spherical Symmetry*, $\mathbf{I} = \text{diag}\{I, I, I\}$. In this case, the standard formulas follow immediately from (43) for $\alpha = \gamma = 1$:

$$\vec{\Omega}_G = \frac{3GM}{2r^3} (\vec{r} \times \vec{V}), \quad \vec{\Omega}_{LT} = \frac{2GI}{r^3} \left[-\vec{\omega} + \frac{3}{r^2} (\vec{\omega} \cdot \vec{r})\vec{r} \right] \tag{44}$$

Note that we have thus shown this to be the exact result for a spherical Earth with any *radial* density distribution $\rho = \rho(r)$.

(b) *Symmetric Top*, $\mathbf{I} = \text{diag}\{I_1, I_1, I\}$, $I_1 \neq I$. It turns out that for $\vec{\omega} = \omega \hat{z}$, i.e., for the rotation about the material symmetry axis, the previous expression for Ω_{LT} remains true; to lowest order in the oblateness, this also proves to be the exact result for a slightly oblate uniform ellipsoid of revolution rotating about its semiminor axis. The corresponding expression for Ω_G is given and discussed in Section 9.

8. EARTH MODELS

To go beyond the $l \leq 2$ approximation, one must make some assumptions about the shape of the central body and density distribution inside it, and use the data of gravitational potential measurements, as available. Bearing in mind the application to GP-B, that is, the Earth, we use the following set of assumptions.

1. *Gravitational Potential*. We assume it to be measured, i.e., the gravitational coefficients a_m^v known. Our problem is thus to determine the vector potential in terms of the coefficients ρ_{lm}^{iv} using as general an Earth model as possible.

2. *Shape*. We assume that the Earth is a slightly oblate ellipsoid of revolution, so its surface equation to lowest order in the eccentricity ϵ is

$$r = r_s(\theta, \varphi) = R(1 - \epsilon \cos^2\theta) \tag{45}$$

where R is the semimajor axis (equatorial radius); the eccentricity is the ratio of the difference of the semiaxes to the major one.

3. *Mass Distribution*. We examine two different models:

(a) With $\rho_0 > 0$ and $\Delta\rho$ being arbitrary functions of their arguments, we set

$$\rho(\vec{r}) = \rho_0(r) + \Delta\rho(\theta, \varphi) > 0, \quad \int_{\text{sphere}} \Delta\rho(\theta, \varphi) \sin\theta \, d\theta \, d\varphi = 0 \tag{46}$$

The first term here describes any depth variation of the *average* density, and the only assumption is that the angular variations are depth independent.

(b) For arbitrary functions $\rho_0 > 0$ and ρ_s , we set

$$\rho(\vec{r}) = \rho_0(r) + \rho_s(\theta, \varphi) \delta(r - r_s(\theta, \varphi)) > 0 \tag{47}$$

Contrary to the previous model, here all angular variations of the density are concentrated at the earth's surface.

$$\vec{\Omega}_G = \frac{(\alpha + 2\gamma)G}{2r^3} \left\{ \left[M + \frac{3}{2r^2} \left(\text{tr } \mathbf{I} - \frac{5}{r^2} (\mathbf{I}\vec{r} \cdot \vec{r}) \right) \right] (\vec{r} \times \vec{V}) + \frac{3}{r^2} (\mathbf{I}\vec{r} \times \vec{V}) \right\} \tag{42}$$

$$\vec{\Omega}_{LT} = \frac{(\alpha + \gamma)G}{r^3} \left\{ \mathbf{I}\vec{\omega} - 3 \left[\frac{1}{2} (\text{tr } \mathbf{I}) - \frac{1}{r^2} (\mathbf{I}\vec{r} \cdot \vec{r}) \right] \vec{\omega} + 3 \frac{\vec{\omega} \cdot \vec{r}}{r^2} \left[\frac{1}{2} (\text{tr } \mathbf{I})\vec{r} - \mathbf{I}\vec{r} \right] \right\} \tag{43}$$

[Of course, the same expression for $\vec{\Omega}_{LT}$ is also obtained from (40) and (39) with $l = 2$.]

Note that formulas (40) and (41) hold in both the body-fixed and inertial frames, and the coefficients and moments of inertia in the latter depend on the time if the rotation is not axisymmetric, $a_{lm}^v = a_{lm}^v(t)$, $p_{lm}^{iv} = p_{lm}^{iv}(t)$, $I_{ij} = I_{ij}(t)$. The expressions (42) and (43) for the precessions are meaningful in the inertial frame, where generally $\mathbf{I} = \mathbf{I}(t) = \mathbf{R}(t)\mathbf{I}(0)\mathbf{R}^T(t)$, $\mathbf{R}(t)$ being a rotation matrix which converts the body-fixed radius vector into the inertial one. In the above case of a simple rotation about an inertially fixed axis, $\vec{\omega} = \omega\hat{z}$,

$$\mathbf{R}(t) = \begin{vmatrix} \cos \omega t & -\sin \omega t & 0 \\ \sin \omega t & \cos \omega t & 0 \\ 0 & 0 & 1 \end{vmatrix}$$

The results (41)–(43) are valid under either of the two conditions (1) *far field*, $R/r \ll 1$; (2) *high symmetry*, i.e., all higher order moments are small. These results alter somewhat our usual notion of the geodetic and Lense–Thirring effects: the first one is generally proportional not only to the orbital momentum, but to $\mathbf{I}\vec{r} \times \vec{V}$ as well, and the Lense–Thirring precession generally points not only in the direction $\vec{\omega}$ of the angular momentum $\vec{L} = \mathbf{I}\vec{\omega}$, but has also components parallel to $\vec{\omega}$, r , and $\mathbf{I}\vec{r}$. Two particular cases of the inertia tensor are of special interest.

(a) *Spherical Symmetry*, $\mathbf{I} = \text{diag}\{I, I, I\}$. In this case, the standard formulas follow immediately from (43) for $\alpha = \gamma = 1$:

$$\vec{\Omega}_G = \frac{3GM}{2r^3} (\vec{r} \times \vec{V}), \quad \vec{\Omega}_{LT} = \frac{2GI}{r^3} \left[-\vec{\omega} + \frac{3}{r^2} (\vec{\omega} \cdot \vec{r})\vec{r} \right] \tag{44}$$

Note that we have thus shown this to be the exact result for a spherical Earth with any *radial* density distribution $\rho = \rho(r)$.

→ (b) *Symmetric Top*, $\mathbf{I} = \text{diag}\{I_1, I_1, I\}$, $I_1 \neq I$. It turns out that for $\vec{\omega} = \omega \hat{z}$, i.e., for the rotation about the material symmetry axis, the previous expression for Ω_{LT} remains true; to lowest order in the oblateness, this also proves to be the exact result for a slightly oblate uniform ellipsoid of revolution rotating about its semiminor axis. The corresponding expression for Ω_G is given and discussed in Section 9.

8. EARTH MODELS

To go beyond the $l \leq 2$ approximation, one must make some assumptions about the shape of the central body and density distribution inside it, and use the data of gravitational potential measurements, as available. Bearing in mind the application to GP-B, that is, the Earth, we use the following set of assumptions.

1. *Gravitational Potential*. We assume it to be measured, i.e., the gravitational coefficients a_m^v known. Our problem is thus to determine the vector potential in terms of the coefficients p_{lm}^v using as general an Earth model as possible.

2. *Shape*. We assume that the Earth is a slightly oblate ellipsoid of revolution, so its surface equation to lowest order in the eccentricity ϵ is

$$r = r_s(\theta, \varphi) = R(1 - \epsilon \cos^2\theta) \tag{45}$$

where R is the semimajor axis (equatorial radius); the eccentricity is the ratio of the difference of the semiaxes to the major one.

3. *Mass Distribution*. We examine two different models:

(a) With $\rho_0 > 0$ and $\Delta\rho$ being arbitrary functions of their arguments, we set

$$\rho(\vec{r}) = \rho_0(r) + \Delta\rho(\theta, \varphi) > 0, \quad \int_{\text{sphere}} \Delta\rho(\theta, \varphi) \sin \theta \, d\theta \, d\varphi = 0 \tag{46}$$

The first term here describes any depth variation of the *average* density, and the only assumption is that the angular variations are depth independent.

(b) For arbitrary functions $\rho_0 > 0$ and ρ_s , we set

$$\rho(\vec{r}) = \rho_0(r) + \rho_s(\theta, \varphi) \delta(r - r_s(\theta, \varphi)) > 0 \tag{47}$$

Contrary to the previous model, here all angular variations of the density are concentrated at the earth's surface.

We will call model A the set of assumptions 1, 2, and 3(a), and model B the assumptions 1, 2, and 3(b); in both cases the mass distributions are, of course, assumed consistent with the gravitational data of the assumption 1.

As far as the Earth is concerned, assumption 2 reflects the classical Clairaut formula (Roy, 1988) (the eccentricity is $\epsilon \approx 3.353 \times 10^{-3}$); also, the mass model 3(b) seems rather realistic for the Earth since the estimated thickness of the layer where its mass distribution varies significantly in the angular directions is only about 30 km.

Note that an entirely different Earth model is used in geodesy: it assumes that the sum of gravitational and centrifugal potentials is constant at the Earth ellipsoid's surface Kaula (1966). This allows one to relate a_{10} to the eccentricity and the Earth angular velocity only (in particular, to obtain the Earth's gravitational oblateness $J_2 = -a_{20}$ with a surprisingly good accuracy), but gives zero values to a_{lm}^v , $m \neq 0$, and leaves p_{lm}^{iv} undetermined.

Evidently, our two models should give bounds for the corrections to the $l \geq 2$ values of the precessions. Moreover, for both of them it is possible to find the corrections explicitly by means of the following four steps:

1. Calculate a_{lm}^v by (32) via spherical harmonics coefficients ρ_{lm}^v of the function $\Delta\rho(\theta, \varphi)$ for model A [or $\rho_s(\theta, \varphi)$ for model B].

2. Since a_{lm}^v are assumed known, solve the resulting equations for ρ_{lm}^v , hence having them expressed via a_{lm}^v (thus $\Delta\rho$ or ρ_s are uniquely determined at this stage through the gravitational data).

3. Using that, calculate the needed moments $M_{l+1l-1m}^v$ of the density by (36) in terms of a_{lm}^v .

4. Using the found values of $M_{l+1l-1m}^v$, express p_{lm}^{iv} through a_{lm}^v according to (38).

In fact, what is described here is a fit of our density distributions (46) and (47) to the known gravitational coefficients a_{lm}^v , which allows us, in these two cases, to express the former through the latter and thus determine uniquely the coefficients p_{lm}^{iv} , i.e., the gravitomagnetic part of the field. The implementation of this procedure is rather cumbersome, though basically straightforward, so the details are given in the Appendix. The results of the calculations to lowest order in the Earth's eccentricity are

$$p_{lm}^{iv} = \frac{1}{2(2l+1)} \left[-(l+m+1)(l+m+2) a_{l+1m+1}^v + (l-m) \right. \\ \left. \times (l-m-1) \kappa_l a_{l-1m+1}^v + \frac{2}{2-\delta_{m1}} (a_{l+1m-1}^v - \kappa_l a_{l-1m-1}^v) \right] \\ + O(\epsilon^2)$$

$$\begin{aligned}
 p_{lm}^{2\nu} &= \frac{(\mp)}{2(2l+1)} \left[(l+m+1)(l+m+2)a_{l+1m+1}^\mu + (l-m) \right. \\
 &\quad \times (l-m-1)\kappa_l a_{l-1m+1}^\mu - \frac{2}{2-\delta_{m1}} (a_{l+1m-1}^\mu + \kappa_l a_{l-1m-1}^\mu) \left. \right] \\
 &\quad + O((\epsilon l)^2) \\
 p_{lm}^{3\nu} &= \frac{1}{2l+1} [(l+m+1)a_{l+1m}^\nu + (l-m)\kappa_l a_{l-1m}^\nu] + O((\epsilon l)^2) \tag{48}
 \end{aligned}$$

with the minus sign and $\mu = s$ for $\nu = c$, the plus sign and $\mu = c$ for $\nu = s$ in the second of these formulas, and

$$\kappa_l = \frac{l+2}{l+4} \quad (\text{model A}), \quad \kappa_l = 1 \quad (\text{model B})$$

It is remarkable that the first nonzero oblateness correction is quadratic, and that the results for the two models differ by just a factor $(l+2)/(l+4)$ in front of a couple of terms. These expressions are needed only for $l \geq 2$ since for $l = 1$ formulas (40) always hold.

9. RESULTS FOR GRAVITY PROBE B

We now apply the above results to the GP-B satellite which will be circling the Earth on a low ($R/r \approx 0.9$) polar, nearly circular orbit. Our aim is to check the GR predictions for the geodetic and Lense–Thirring drift of a GP-B gyroscope taking into account all the peculiarities of the real Earth’s gravitoelectric and gravitomagnetic field pertinent to the expected experimental error of 1 part in 10^5 for the geodetic effect and about 1 part in 10^2 for the Lense–Thirring effect. More precisely, one needs to check whether the spherically symmetric approximation (44), which has been used for many years in the theoretical discussion and planning of the experiment, needs any corrections to match the above experimental accuracy.

We use the GR values for the Eddington parameters $\alpha = \gamma = 1$ in the sequel. We always use assumption 2 of the previous section that the Earth has a shape of a slightly oblate ellipsoid of revolution with the eccentricity $\epsilon = 3.353 \times 10^{-3}$. Its semiminor axis is assumed fixed in the inertial space and coincident with the Earth rotation axis and the z axis of the Cartesian coordinates.

1. Gravitational Potential. For the required accuracy, it is enough to include only the Earth’s quadrupole moment into Φ , i.e., to set

$$-a_{20} \equiv J_2 \approx 1.083 \times 10^{-3}; \quad a_{lm}^v = 0 \quad \text{for } l = 2, m > 0; \quad l > 2 \tag{49}$$

This is because all gravitational coefficients other than the Earth’s oblateness J_2 are at least 2 orders of magnitude smaller (*World Geodetic System 1984, 1987*). By (49) and (40), the $l \leq 2$ expression (40) for Φ is valid with

$$\mathbf{I} = \text{diag}\{I_1, I_1, I\}, \quad I_1 = I - J_2MR^2 \tag{50}$$

which gives the gravitational potential in the usual form:

$$\Phi(\vec{r}) = -\frac{GM}{r} \left[1 - J_2 \frac{R^2}{2r^2} \left(3 \frac{z^2}{r^2} - 1 \right) \right] = -\frac{GM}{r} \left[1 - J_2 \frac{R^2}{r^2} P_2(\cos \theta) \right]$$

2. *Instantaneous Geodetic Precession.* Just as for the potential, (49) assures that the $l \leq 2$ formula (42) is valid for Ω_G . The explicit form of the geodetic precession is obtained either from (42) and (50) or from its general definition (18) by differentiating the above expression for the potential. The result reads

$$\begin{aligned} \vec{\Omega}_G = \frac{3GM}{2r^3} & \left\{ (\vec{r} \times \vec{V}) + J_2 \frac{3}{2} \left(\frac{R}{r} \right)^2 \right. \\ & \left. \times \left[\left(1 - 5 \frac{z^2}{r^2} \right) (\vec{r} \times \vec{V}) + 2z(\hat{z} \times \vec{V}) \right] \right\} \end{aligned} \tag{51}$$

This expression contains the J_2 corrections to the classical spherically symmetric expression (44). Breakwell (1988) obtained it, but in a different form, pinning the value of the precession to a point on a given satellite orbit rather than to a given point in space through which the spinning particle passes with a velocity \vec{V} .

3. *Gravitomagnetic Field and the Instantaneous Lense–Thirring Precession.* We now invoke our assumption about the mass distribution (models A and B of the previous section) and substitute the values (49) of the gravitational coefficients into the expressions (48) for p_{lm}^i . In this way it turns out that for both models A and B the only nonzero coefficients additional to $l = 1$ are those with $l = 3, m = 0, 1$, and their values are

$$\begin{aligned} p_{31}^{1c} = p_{31}^{2s} = \frac{5}{49} J_2, \quad p_{30}^3 = -\frac{15}{49} J_2 & \quad (\text{model A}) \\ p_{31}^{1c} = p_{31}^{2s} = \frac{1}{7} J_2, \quad p_{30}^3 = -\frac{3}{7} J_2 & \quad (\text{model B}) \end{aligned} \tag{52}$$

So, for both our models the only addition to the field of a symmetric top is the gravitomagnetic field (52). In terms of the Lense–Thirring precession that means that the standard expression (44) has to be appended by the terms (52) from the general formula (39) for the precession. The result reads

$$\begin{aligned} \vec{\Omega}_{LT} = & \frac{GI\omega}{r^3} \left[\left(3 \frac{z^2}{r^2} - 1 \right) \hat{z} + 3 \frac{z}{r} \frac{x\hat{x} + y\hat{y}}{r} \right] \\ & - J_2 \frac{GMR^2\omega}{r^3} \frac{R}{r} \left[Z \left(\frac{z^2}{r^2} \right) \frac{R}{r} \hat{z} + X \left(\frac{z^2}{r^2} \right) \frac{x\hat{x} + y\hat{y}}{r} \right] \end{aligned} \quad (53)$$

where (recall $z^2/r^2 = \cos^2\theta$)

$$\begin{aligned} Z(\cos^2\theta) &= Z_0(35 \cos^4\theta - 30 \cos^2\theta + 3), & X(\cos^2\theta) &= X_0(5 \cos^2\theta - 1) \\ Z_0 &= \frac{15}{98}, & X_0 &= \frac{15}{49} \quad (\text{model A}); \\ Z_0 &= \frac{3}{14}, & X_0 &= \frac{3}{7} \quad (\text{model B}) \end{aligned} \quad (54)$$

The second line of (53) is the correction to the classical Lense–Thirring expression in the first line which is induced by the gravitomagnetic field (52). [Recall from Section 7, case (b), that the dependence of $\vec{\Omega}_{LT}$ on J_2 for a symmetric top rotating about its symmetry axis cancels out—remarkably! Thus the first line of (53) contains no J_2 .]

Note that the only difference between the Earth models A and B is a factor of 7/5 in the constants Z_0 and X_0 .

Note also that allowing for more multipoles in the gravitational potential than given in equation (49) automatically yields higher order multipoles in the gravitomagnetic field by the expressions (48) for p_{lm}^i . However, for GP-B all these higher order corrections prove to be too small to be taken into account.

4. Orbit of the GP-B Satellite and the Instantaneous Values of Precessions. The perfect orbit for the GP-B satellite would be a circular polar one with the altitude $h_s = 650$ km, which is described by $r = r_0 = R + h_s$, $V = V_0 = \sqrt{GM/r_0}$. In reality it is slightly distorted by the quadrupole moment of the Earth's gravitational field; with the lowest order in J_2 corrections included, the orbit becomes (Breakwell, 1987; Barker and O'Connell, 1970)

$$r = r_0 \left[1 - \frac{1}{4} J_2 \left(\frac{R}{r_0} \right)^2 \cos 2\theta \right]$$

$$\vec{V} = V_0 \left[1 - \frac{3}{8} J_2 \left(\frac{R}{r_0} \right)^2 \right] \left[\hat{\theta} + \frac{1}{2} J_2 \left(\frac{R}{r_0} \right)^2 \sin 2\theta \hat{r} \right]$$
(55)

where \hat{r} , $\hat{\theta}$, $\hat{\phi}$ are the corresponding unit vectors. (In view of the orbit symmetry, it is enough to consider a half-orbit from one pole to the other, $0 \leq \theta \leq \pi$; otherwise, the second half may be assigned to $\pi \leq \theta \leq 2\pi$, with the proper direction of the unit vector \hat{e}_θ .)

Introducing the orbital velocity (55) into the formula (51) and dropping some terms $O(J_2^2)$, we obtain the geodetic precession for the GP-B gyroscope in the following form:

$$\vec{\Omega}_G = \frac{3GMV_0}{2r^2} \left[1 - \frac{3}{8} J_2 \left(\frac{R}{r_0} \right)^2 \right] \left[1 + \frac{3}{2} J_2 \left(\frac{R}{r_0} \right)^2 (1 - 3 \cos^2 \theta) \right] \hat{e}_\phi$$

Using now the orbit radius (55) and keeping only the corrections linear in J_2 , we arrive at the expression we need:

$$\vec{\Omega}_G = \frac{3GMV_0}{2r_0^2} \left[1 + J_2 \left(\frac{R}{r_0} \right)^2 \left(\frac{5}{8} - \frac{7}{2} \cos^2 \theta \right) \right] \hat{e}_\phi$$
(56)

To derive a similar formula for $\vec{\Omega}_{LT}$ from (53), we of course need no orbital velocity, but only the orbit radius (55); with accuracy $O(J_2)$ the instantaneous value of the Lense–Thirring precession for GP-B turns out to be

$$\vec{\Omega}_{LT} = \vec{\Omega}_{LT}^{(0)} + \vec{\Omega}_{LT}^{(J_2)}$$

$$\vec{\Omega}_{LT}^{(0)} = \frac{GI\omega}{r_0^3} [(3 \cos^2 \theta - 1)\hat{z} + (3 \cos \theta \sin \theta)\hat{x}]$$
(57)

$$\vec{\Omega}_{LT}^{(J_2)} = J_2 \frac{GI\omega}{r_0^3} \left[\left(\frac{R}{r_0} \right)^2 \left(U - Z \frac{MR^2}{I} \right) \hat{z} + \left(V \frac{R}{r_0} \cos \theta - X \frac{MR^2}{I} \right) \sin \theta \hat{x} \right]$$

For convenience, we have assumed that the orbit is in the plane $y = 0$; the values of $Z = Z(\cos^2 \theta)$ and $X = X(\cos^2 \theta)$ are given in (54), while

$$U(\cos^2 \theta) = \frac{3}{4} (6 \cos^4 \theta - 5 \cos^2 \theta + 1), \quad V(\cos^2 \theta) = \frac{9}{4} (2 \cos^2 \theta - 1)$$
(58)

Note that the expressions (56) and (57) contain the corrections coming both directly from the non-spherically symmetric gravitoelectric and gravito-

magnetic field and also through the influence of the former on the orbit. The first contribution can be distinguished from the second in the Lense–Thirring precession (57) by the presence of the factor MR^2/I ; however, in the geodetic precession (56) this difference is concealed.

5. *Orbit-Averaged Precessions: the GP-B Drift Rates.* With the above results, it is now straightforward to carry out the orbit averaging, which reduces to a simple integration over θ . In this way, to lowest order in the Earth's oblateness $J_2 \approx 1.083 \times 10^{-3}$, using (56), we find

$$\langle \vec{\Omega}_G \rangle = \frac{3GMV}{2r_0^2} \left[1 - \frac{9}{8} J_2 \left(\frac{R}{r_0} \right)^2 \right] \hat{e}_\varphi = \frac{3GMV}{2r_0^2} [1 - 1.003 \times 10^{-3}] \hat{e}_\varphi \quad (59)$$

where the values $R = 6378$ km and $r_0 = 7028$ km for the Earth's and orbit radius have been taken. The result exactly coincides with the one found by Breakwell (1988) in a different way (he gave the formula for the orbit of any inclination). Since GP-B is intended to measure the geodetic precession and the Eddington parameter γ to about 1 part in 10^5 , a 0.1% correction is critically important. Note that the GP-B gyro spin axis will be initially aligned with the reference direction to the guide star in the orbit plane, so the geodetic effect will cause the in-plane drift of the spin in the north–south direction.

In a similar fashion, from (57) we derive

$$\langle \vec{\Omega}_{LT} \rangle = \frac{GI\omega}{2r_0^3} \left[1 + \frac{9}{4} J_2 \left(\frac{R}{r_0} \right)^2 \left(\frac{1}{2} - Z_0 \frac{MR^2}{I} \right) \right] \hat{z} \quad (60)$$

For our model B, $Z_0 = 3/14$ according to (54); therefore, with $MR^2/I = 3.024$ for the Earth,

$$\langle \vec{\Omega}_{LT} \rangle = \frac{GI\omega}{2r_0^3} \left[1 + \frac{9}{8} J_2 \left(\frac{R}{r_0} \right)^2 \left(1 - \frac{3}{7} \frac{MR^2}{I} \right) \right] \hat{z} = \frac{GI\omega}{2r_0^3} [1 - 2.97 \times 10^{-4}] \hat{z}$$

Thus the J_2 correction for model B is (-0.03%), which is essentially beyond the expected GP-B accuracy for the Lense–Thirring effect. For model A the correction is of the opposite sign and almost one order of magnitude smaller, about 0.007%. For a different model of the mass distribution inside the Earth, Teyssandier (1977) obtained a slightly larger correction of about -0.011% , with the same significance for GP-B.

6. *Effect of the Moon.* According to formula (21) from Section 5, the geodetic effect from a distant mass such as the Moon scales with the mass and inversely with the square of distance. Since the Moon has a small mass

and is at a large distance, we expect its effect to be small. A rough estimate for the geodetic precession due to the Moon is

$$\frac{\Omega_G^M}{\langle \Omega_G \rangle} = \frac{M_M}{M} \left(\frac{r_0}{r_M} \right)^2 \approx 10^{-6} \quad (62)$$

which is too small to be of significance for the GP-B experiment.

The Lense–Thirring effect scales with the angular momentum and inversely with the cube of the distance. Since the velocity of the Moon in its orbit is small, we expect its effect to be small, and the estimate

$$\frac{\Omega_{LT}^M}{\langle \Omega_{LT} \rangle} = \frac{L_M}{L} \left(\frac{r_0}{r_M} \right)^3 \approx 10^{-5} \quad (63)$$

shows that it is really far beyond the GP-B accuracy. Thus the Moon is of no consequence for the GP-B experiment.

6. *Effect of the Sun.* Although the Sun is quite distant, its mass is large compared to the Earth, so it is not apparent how large its effect on the precession will be. We can calculate this using (18). In the inertial frame centered on the Sun the velocity of the satellite will be the sum of the satellite velocity in the Earth orbit plus the orbital velocity of the Earth,

$$\vec{V} = \vec{V}_S + \vec{V}_{EO} \quad (64)$$

But the satellite velocity averages to zero in the course of one orbit, while the other factors in (18) change very little, so for long time averages we may neglect the satellite velocity compared to the Earth orbital velocity. Treating the Sun as a point mass and approximating the Earth orbit as a circle, we may thus write (18) as

$$\vec{\Omega}_G^S \approx \left(\frac{\alpha + 2\gamma}{2} \right) \frac{GM_S V_{EO}}{r_S^2} \hat{n} \quad (65)$$

where the unit vector \hat{n} is perpendicular to the plane of the ecliptic and r_S is the Earth–Sun distance. The numerical value of this, with $\alpha = \gamma = 1$, is

$$\langle \Omega_G^S \rangle \approx 6.3 \times 10^{-7} \text{ rad/year} = 19 \text{ marcsec/year} \quad (66)$$

This contribution to the GP-B precession is not negligible and must be included in the data analysis. It was first discussed by deSitter (1916a, b), and further information and references can be found in the book by Will (1993). Note an important fact concerning the direction of the above precession vector: it is perpendicular to the plane of the ecliptic, whereas the geodetic precession vector due to the Earth lies mainly in the equatorial plane of the Earth, so the two are not parallel. Indeed, the precession due to the

Sun is roughly in the direction of the Lense–Thirring precession due to the Earth. Finally, we may estimate the precession due to the spin of the Sun by using (44). Since the spin period of the Sun is about 1 month, this gives roughly

$$\langle \Omega_{LT}^S \rangle \approx 10^{-11} \text{ rad/year} = 0.002 \text{ marcsec/year} \quad (67)$$

which is far too small to be of relevance to the GP-B experiment.

ACKNOWLEDGMENTS

This work was supported by NASA grant NAS 8-39225 to Gravity Probe B. We are grateful to the members of GP-B Theory Group, especially, to C. W. F. Everitt, G. M. Keiser, R. V. Wagoner, and P. W. Worden, for many fruitful discussions and enlightening comments.

APPENDIX

Let us show briefly the implementation of the four-step procedure described in Section 8, which allows us to obtain expressions (48) for gravitomagnetic multipole coefficients p_{lm}^v . We do it for the Earth model A; calculations for model B are similar.

First, we introduce the model A density distribution (46) into the definition (36) of the general moment M_{klm}^v of the density and calculate this moment for the shape of a slightly oblate ellipsoid of revolution (assumption 2 of Section 8). Working to the first order in the eccentricity ϵ , we find M_{klm}^v in terms of the spherical harmonics coefficients

$$\rho_{lm}^v = \int \Delta \rho Y_{lm}^v \sin \theta \, d\theta \, d\varphi$$

of the function $\Delta \rho(\theta, \varphi)$ from (46), namely

$$R^{-3} M_{klm}^v = \frac{1 - \epsilon(k+3)Q_{lm}}{k+3} \rho_{lm}^v - \epsilon(k+3)[S_{lm}\rho_{l+2,m}^v + T_{lm}\rho_{l-2,m}^v] + O((\epsilon k)^2) \quad (\text{A.1})$$

Here Q_{lm} , S_{lm} , T_{lm} are known positive rational fractions of l and m bounded for all pertinent values of those parameters,

$$Q_{lm} = \frac{(l-m+1)(l+m+1)}{(2l+1)(2l+3)} + \frac{(l+m)(l-m)}{(2l-1)(2l+1)}$$

$$S_{lm} = \frac{(l-m+1)(l-m+2)}{(2l+1)(2l+3)}, \quad T_{lm} = \frac{(l+m)(l+m-1)}{(2l-1)(2l+1)}$$

Formula (A.1) is slightly different for the case $l = m = 0$, but we do not need it.

For $k = 1$ the left-hand side of the equality (A.1) is given via a_{lm}^y according to (37):

$$R^{-3} \frac{M}{2 - \delta_{m0}} \frac{(l + m)!}{(l - m)!} a_{lm}^y = \frac{1 - \epsilon(l + 3)Q_{lm}}{l + 3} \rho_{lm}^y - \epsilon(l + 3)[S_{lm}\rho_{l+2,m}^y + T_{lm}\rho_{l-2,m}^y] + O((\epsilon l)^2)$$

This is a tridiagonal system of linear algebraic equations for ρ_{lm}^y with small off-diagonal elements. Solving it for ρ_{lm}^y , we express the latter in terms of a_{lm}^y :

$$\begin{aligned} \frac{R^3}{M} \rho_{lm}^y &= \frac{l + 3}{N_{lm}} a_{lm}^y \\ &+ \epsilon \left[\frac{l + 3}{N_{lm}} Q_{lm} a_{lm}^y + \frac{l + 5}{N_{l+2m}} S_{lm} a_{l+2m}^y + \frac{l + 1}{N_{l-2m}} T_{lm} a_{l-2m}^y \right] \\ &+ O((\epsilon l)^2) \end{aligned} \tag{A.2}$$

where

$$N_{lm} = (2 - \delta_{m0}) \frac{(l - m)!}{(l + m)!}$$

Introducing now the expression (A.2) back into the general formula (A.1), we get *all* moments, with *any* k , expressed through the gravitational coefficients:

$$\begin{aligned} M_{klm}^y &= \frac{M}{2 - \delta_{m0}} \left\{ \frac{(l + m)!}{(l - m)!} \frac{l + 3}{k + 3} a_{lm}^y \right. \\ &\left. - \epsilon \frac{k + 2}{k + 3} [\tilde{V}_{lm} a_{lm}^y + \tilde{S}_{lm} a_{l+2,m}^y + \tilde{T}_{lm} a_{l-2,m}^y] \right\} + O((\epsilon l)^2) \end{aligned} \tag{A.3}$$

with the quantities \tilde{Q}_{lm} , \tilde{S}_{lm} , \tilde{T}_{lm} simply related to Q_{lm} , S_{lm} , T_{lm} , respectively:

$$\begin{aligned} \tilde{Q}_{lm} &= \frac{(l + m)!}{(l - m)!} Q_{lm}, & \tilde{S}_{lm} &= \frac{(l + m + 2)!}{(l - m + 2)!} S_{lm}, \\ \tilde{T}_{lm} &= \frac{(l + m - 2)!}{(l - m - 2)!} T_{lm} \end{aligned}$$

We finally set $k = l \pm 1$ in (A.3) and use the resulting expressions in the

relation (38), completing thus the last step of the procedure described in Section 8 and obtaining the equalities (48) for p_{lm}^{iv} (model A).

REFERENCES

- Adler, R. J. (1999). *Gen. Rel. Grav.* **31**, 1837.
- Adler, R. J., and Silbergleit, A. S. (1998). In *Proceedings of the IVth Alexander Friedmann International Seminar on Gravitation and Cosmology*, St. Petersburg, Russia, June 17–25, 1998. Eds. Yu. N. Gnedin, A. A. Grib, V. M. Mostepanenko, W. A. Rodrigues, Jr., pp. 102–115.
- Barker, B. M., and O'Connell, R. F. (1970). *Phys. Rev.* **D2**, 1428.
- Bateman, H., and Erdélyi, A. (1953). *Higher Transcendental Functions*, Vol. 1, McGraw-Hill New York, sec. 3.9.
- Breakwell, J. V. (1987). Stanford University Class Notes, Advanced Space Mechanics AA279B, Stanford, California.
- Breakwell, J. V., (1988). In *Near Zero*, J. D. Fairbank, B. S. Deaver, Jr., C. W. F. Everitt, and P. F. Michelson, eds., Freeman, New York, pp. 685–690.
- Bursa, M. (1992). *Studia Geophys. Geodes.* **36**, 109.
- Ciufolini, I., and Wheeler, J. A. (1995). *Gravitation and Inertia*, Princeton University Press, Princeton, New Jersey, Chapter 6.
- de Sitter, W. 1916a, MNRAS 76, 699.
- de Sitter, W. 1916b, MNRAS 77, 155.
- Eddington, A. S. (1988). *The Mathematical Theory of Relativity*, Cambridge University Press, New York, p. 105.
- Kaula, W. M. (1966). *Theory of Satellite Geodesy*, Blaisdell, Waltham, Massachusetts.
- Keiser, G. M. (1998). In *Proceedings of the 3rd W. Fairbank Meeting, Rome—Pescara*, June 29–July 4, in press.
- Lense, J., and Thirring, H. (1918). *Phys. Z.* **19**, 156.
- Misner, C. W., Thorne, K. S., and Wheeler, J. A., (1973). *Gravitation*, Freeman, New York, sec. 40.7.
- Nordtvedt, K. (1968). *Phys. Rev. D.* **169**, 1017.
- Nordtvedt, N. (1988a). *Int. J. Theor. Phys.* **27**, 1395.
- Nordtvedt, K. (1988b). *Phys. Rev. Lett.* **61**, 2647.
- Nordtvedt, K. (1998). In *Proceedings of the 3rd W. Fairbank Meeting, Rome—Pescara*, June 29–July 4, in press.
- Ohanian, H. C., and Ruffini, R. (1994). *Gravitation and Spacetime*, Norton, New York.
- Robertson, H. P. (1962). In *Space Age Astronomy*, A. J. Deutsch and W. B. Klemperer, eds., Academic Press, New York, p. 128.
- Rose, E. M. (1957). *Elementary Theory of Angular Momentum*, Wiley, New York.
- Roy, A. E. (1988). *Orbital Motion*, Adam Hilger, Bristol, England.
- Teyssandier, P. (1977). *Phys. Rev. D* **16**, 946.
- Teyssandier, P. (1978). *Phys. Rev. D* **18**, 1037.
- Wilkins, D. C. (1968). *Ann. Phys. (NY)* **61**, 277.
- Will, C. M. (1993). *Theory and Experiment in Gravitational Physics*, Cambridge University Press, Cambridge.
- World Geodetic System 1984*. (1987). DMA Technical Report.